

In vivo incorporation of multiple unnatural amino acids through nonsense and frameshift suppression

Erik A. Rodriguez*, Henry A. Lester†, and Dennis A. Dougherty**

Divisions of *Chemistry and Chemical Engineering and †Biology, California Institute of Technology, Pasadena, CA 91125

Edited by Peter G. Schultz, The Scripps Research Institute, La Jolla, CA, and approved April 21, 2006 (received for review December 15, 2005)

Site-specific incorporation of unnatural amino acids (UAAs) into proteins is a valuable tool for studying structure–function relationships, incorporating biophysical probes, and elucidating protein–protein interactions. In higher eukaryotic cells, the methodology is currently limited to incorporation of a single UAA in response to a stop codon, which is known as nonsense suppression. Frameshift suppression is a unique methodology for incorporating UAAs in response to quadruplet codons, but currently, it is mostly limited to *in vitro* protein translation systems. Here, we evaluate the viability of frameshift suppression in *Xenopus* oocytes. We demonstrate UAA incorporation by using yeast phenylalanine frameshift suppressor (YFFS) tRNAs that recognize two different quadruplet codons (CGGG and GGGU) *in vivo*. Suppression efficiency of the YFFS tRNAs increases nonlinearly with the amount of injected tRNA, suggesting a significant competition with endogenous, triplet-recognizing tRNA. Both frameshift suppressor tRNAs are less efficient than the amber suppressor tRNA THG73 (*Tetrahymena thermophila* G73), which has been used extensively for UAA incorporation in *Xenopus* oocytes. However, the two YFFS tRNAs are more “orthogonal” to the *Xenopus* system than THG73, and they offer a viable replacement when suppressing at promiscuous sites. To illustrate the potential of combining nonsense and frameshift suppression, we have site-specifically incorporated two and three UAAs simultaneously into a neuroreceptor expressed *in vivo*.

nicotinic receptor | tRNA | quadruplet codon | stop codon | protein engineering

The site-specific incorporation of unnatural amino acids (UAAs) into proteins biosynthetically is a powerful methodology that is seeing increasing use. The primary approach has been stop codon (nonsense) suppression using a specially designed tRNA with an anticodon that recognizes the stop codon. A wide range of *in vitro* translation systems has been used, along with expression in *Escherichia coli* and, to a lesser extent, yeast. Nonsense suppression in higher eukaryotes has, for the most part, been limited to the *Xenopus* oocyte, where microinjection of the required mRNA and aminoacyl tRNA is straightforward and electrophysiology provides a sensitive probe of UAA incorporation (1, 2). Other experiments in higher eukaryotes have relied on the evolution of a unique tRNA and a complementary aminoacyl-tRNA synthetase (aaRS) to insert a UAA in response to the UAG or UGA stop codon, but currently, only 3-iodo-tyrosine (3), *p*-benzoyl-phenylalanine (4), and 5-hydroxy-tryptophan (5) have been incorporated.

A remarkable variant of this approach is the use of quadruplet codons, a process that is termed frameshift suppression and was pioneered by Sisido and coworkers (6, 7). The success of this approach opens up the possibility of developing multiple additional codons, thus incorporating several different UAAs into a protein. This multiple incorporation, in turn, would enable the use of innovative biophysical approaches such as incorporating FRET pairs, structural probes such as unique cross-linking approaches, and more detailed structure–function studies.

To date, frameshift suppression *in vivo* has been performed only in *E. coli* by using a unique tRNA/aaRS pair, and homoglutamine is the only UAA incorporated by this method. Frameshift suppres-

sion was used simultaneously with nonsense suppression to incorporate two UAAs in *E. coli* (8). It has yet to be established whether frameshift suppression by chemically aminoacylated tRNA can be effective *in vivo* in general and in eukaryotic cells such as the *Xenopus* oocyte in particular. In fact, a previous attempt to perform frameshift suppression in *Xenopus* oocytes showed very poor suppression efficiency (9). Here, we show that with appropriately designed frameshift suppressor (FS) tRNAs, frameshift suppression is a viable approach to UAA incorporation in eukaryotic cells. Also, the efficiency of frameshift suppression can be substantially improved by “masking” the mRNA of all in-frame quadruplet sequences that match the frameshift suppression site. In particular, we describe two tRNAs with four-base anticodons that can deliver UAAs in response to the quadruplet codons CGGG and GGGU. When directly compared with an amber suppressor (AS) tRNA (THG73, *Tetrahymena thermophila* G73) that has been used extensively in *Xenopus* oocytes, the FS tRNAs are less efficient at delivering UAAs. However, both FS tRNAs are more “orthogonal” than THG73, producing much less incorporation of undesired natural amino acids at promiscuous sites. We also show that suppression by FS tRNAs increases nonlinearly with the amount of injected tRNA. To illustrate the potential of this methodology, we have successfully incorporated two and three different UAAs simultaneously into a neuroreceptor expressed in a *Xenopus* oocyte.

Results

Testing Frameshift Suppression Viability *in Vivo*. To determine whether frameshift suppression is viable in *Xenopus* oocytes, we chose to use a tRNA that can be aminoacylated *in vivo*. We selected the human serine AS (HSAS), because it is aminoacylated (with serine) in eukaryotic cells and the seryl-tRNA synthetase does not recognize the anticodon (10–12). The CUA anticodon of HSAS was replaced with CCCG and ACCC to create the human serine FSs (HSFS_{CCCG} and HSFS_{ACCC}) (cloverleaf structures shown in Fig. 1A), which recognize the quadruplet codons CGGG and GGGU. Prior research showed that these four-base codons are efficient *in vitro* (7). Injection of wild-type muscle nicotinic acetylcholine receptor (nAChR) mRNA and either HSFS_{CCCG} or HSFS_{ACCC} (2.5 or 10 ng per oocyte; no amino acid ligated to the tRNA) into *Xenopus* oocytes resulted in no detectable channel expression. The addition of the original AS HSAS with wild-type nAChR mRNA did show channel expression with 2.5 ng of tRNA per oocyte but not with 10 ng. These results suggested that the HSFS tRNAs were causing +1 frameshifts, resulting in undesirable truncation of wild-type protein and thus a lack of detectable current. Analysis of the four nAChR subunits revealed four CGGG and one GGGU in-frame quadruplet codons, which were mutated to degenerate codons (see *Materials and Methods*) to avoid sup-

Conflict of interest statement: No conflicts declared.

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: UAA, unnatural amino acid; aaRS, aminoacyl-tRNA synthetase; AS, amber suppressor; FS, frameshift suppressor; HSAS, human serine AS; HSFS, human serine FS; YFFS, yeast phenylalanine FS; WF1, 5-fluoro-tryptophan; Aba, α -aminobutyric acid; Nval, norvaline; ACh, acetylcholine; nAChR, nicotinic ACh receptor; dCA, dinucleotide deoxyCA.

†To whom correspondence should be addressed. E-mail: dadougherty@caltech.edu.

© 2006 by The National Academy of Sciences of the USA




Fig. 1. tRNAs and UAAs. (A) The AS tRNAs are shown, with the CUA anticodon and the FS anticodons shown at the bottom. YFaFS tRNA acceptor stem mutations are shown next to the YFFS tRNA body (italicized). (B) The three UAAs used in this study.

pression. We refer to the resulting mRNAs as the “masked” constructs. Other groups have similarly removed undesired in-frame quadruplets (7, 9, 13). Injection of 2.5 ng per oocyte of either unligated HSFS plus the masked nAChR mRNAs resulted in functional channels with the same EC_{50} as channels expressed without tRNA (data not shown). Unless otherwise noted, all subsequent experiments used such masked constructs.

To test whether a naturally occurring amino acid (serine) could be incorporated in response to a quadruplet codon, we probed a highly conserved leucine of the nAChR M2 domain, a site designated Leu-9'. This site is a promiscuous site in the nicotinic receptor, and replacement of the native leucine with essentially any natural amino acid produces a functional receptor, usually with a quite noticeable shift in EC_{50} . In particular, prior research showed that a leucine-to-serine mutation in the β -subunit (β 9') resulted in an \approx 33-fold increased sensitivity to acetylcholine (ACh) (14). This site was mutated to UAG, CGGG, or GGGU. When mutant mRNA was injected into *Xenopus* oocytes along with 2.5 ng of unligated HSAS or HSFS tRNA, which should be aminoacylated with serine by the endogenous seryl-tRNA synthetase, significant channel expression was seen. However, the EC_{50} values varied depending on the incubation time (Table 1). This finding suggested that natural amino acids other than serine were being placed at the β 9' site with 2-day incubations, because the conventional mutant, β 9'Ser, shows no change in EC_{50} (Table 1). The variability in EC_{50} between 1- and 2-day incubations suggests that the tRNAs are being modified to accept other amino acids. Modification of yeast phe-

nylalanine tRNA in *Xenopus* oocytes has been shown to increase greatly from 1- to 2- day incubation times (15). Thus, we avoid this complication by incubation for 1 day. Amber suppression is highly efficient when the average maximal peak current (I_{max}) is measured at 1.25 ng of tRNA per oocyte and decreases slightly when 2.5 ng is added (Table 2). CGGG shows lower suppression than GGGU, in agreement with previous *in vitro* studies (7, 16). CGGG suppression is highly nonlinear, with a 330% increase in current when twice as much tRNA is injected (Table 2). GGGU, however, shows an almost linear relationship, with an increase of 86% in response to doubling (Table 2). These data suggest that $\text{HSFS}_{\text{ACCC}}$ is a more efficient tRNA at recognizing its cognate quadruplet codon and/or has less competition with endogenous triplet tRNA in *Xenopus* oocytes than $\text{HSFS}_{\text{CCCG}}$. These experiments establish that frameshift suppression is viable in *Xenopus* oocytes and that UAA incorporation should be feasible when using the appropriate FS tRNA.

UAA Incorporation by Frameshift Suppression. THG73 is an AS tRNA (cloverleaf structure shown in Fig. 1A) (17) that is used extensively for incorporating UAAs into various ion channels expressed in *Xenopus* oocytes (2). Initially, a FS derived from THG73 that recognizes the quadruplet codon CGGG ($\text{THG73FS}_{\text{CCCG}}$) was tested for UAA incorporation. Attempts to suppress β 9'CGGG with $\text{THG73FS}_{\text{CCCG-L}}$, where L was chemically aminoacylated onto the tRNA, showed no current *in vivo*. This result is consistent with data from Voss and coworkers (9), who saw

Table 1. HSAS and HSFS suppression experiments at the β 9' site

β 9'X	tRNA, 2.5 ng	EC_{50} , 1 day*	n_{H}	n	EC_{50} , 2 days*	n_{H}	n
AGC (serine)	none	1.5 ± 0.04	1.7 ± 0.07	5	1.5 ± 0.2	1.9 ± 0.3	3
UAG	HSAS	1.7 ± 0.06	1.7 ± 0.09	6	0.70 ± 0.008	1.9 ± 0.07	14
CGGG	$\text{HSFS}_{\text{CCCG}}$	2.1 ± 0.09	1.7 ± 0.1	8	1.3 ± 0.1	1.9 ± 0.3	13
GGGU	$\text{HSFS}_{\text{ACCC}}$	1.9 ± 0.08	1.5 ± 0.08	9	0.68 ± 0.1	1.7 ± 0.04	5

*Incubation time.

Table 2. HSAS and HSFS suppression experiments at the $\beta 9'$ site (1-day incubation)

$\beta 9'$ X	tRNA	$I_{max} \pm SE,^* 1.25^\dagger$	n	$I_{max} \pm SE,^* 2.5^\dagger$	n	% HSAS, 1.25 [†]	% HSAS, 2.5 [†]	% change [‡]
UAG	HSAS	-19 \pm 2	12	-14 \pm 3	11	100	100	-26
CGGG	HSFS _{CCCG}	-1.3 \pm 0.3	10	-5.6 \pm 1	12	6.8	40	330
GGGU	HSFS _{ACCC}	-8.6 \pm 3	10	-16 \pm 3	12	45	110	86

*Average I_{max} (in μA) recorded at 50 μM ACh.

[†]Nanograms of tRNA.

[‡]Between 1.25 and 2.5 ng.

very little UAA incorporation with THG73FS_{ACCC} in *Xenopus* oocytes.

We then chose to screen yeast phenylalanine FS (YFFS) tRNA, which was used successfully by Sisido and colleagues (7, 16) *in vitro*. We studied both YFFS_{CCCG} and YFaFS_{ACCC} (yeast phenylalanine containing acceptor stem mutations FS); Fig. 1A shows cloverleaf structures. The latter contains acceptor stem mutations (denoted by the "a") incorporated to reduce glycyl-tRNA synthetase recognition (7). We first evaluated a nonpromiscuous position of the nAChR, $\alpha 149W$, an agonist-binding site tryptophan that makes a cation- π interaction with ACh (18). Wild-type recovery (i.e., suppressing the $\alpha 149$ quadruplet codons with YFFS_{CCCG}-W or YFaFS_{ACCC}-W) resulted in functional, wild-type channels (Table 3). To demonstrate UAA incorporation, we relied on previous work using the AS THG73 that established that 5-fluoro-tryptophan (WF1) (structure shown in Fig. 1B) incorporated at $\alpha 149$ decreased the cation- π interaction and caused an ≈ 4 -fold increase in EC₅₀ (18). YFFS_{CCCG}-WF1 suppression at $\alpha 149CGGG$ resulted in a comparable increase in EC₅₀ (Table 3), establishing the successful incorporation of the UAA WF1.

We next considered the previously mentioned Leu-9' residue. Suppression at $\beta 9'$ GGGU and $\delta 9'$ GGGU with YFaFS_{ACCC}-Aba (where Aba is α -aminobutyric acid) and YFaFS_{ACCC}-Nval (where Nval is norvaline) (UAA structures shown in Fig. 1B), respectively, resulted in reductions in EC₅₀ (Table 3) that were consistent with previous studies that used the same UAAs and the AS THG73 (14). All frameshift suppression experiments had an I_{max} between -1.6 and -4.4 μA , which is more than adequate for UAA studies *in vivo* and should allow for the incorporation of multiple UAAs. In all cases, injection of full-length tRNA that had no amino acid attached to the 3' end resulted in no detectable currents in response to added ACh, directly showing a lack of aminoacylation by endogenous *Xenopus* aaRSs.

Masking Effects on Frameshift Suppression. Experiments with HSFS required the masking of the nAChR subunits to avoid protein truncation caused by +1 frameshifts. To demonstrate the effect on UAA incorporation, suppression experiments were performed with wild-type and masked constructs. The quadruplet codon GGGU was chosen because there was only one in-frame quadruplet in the signaling sequence of the nAChR β -subunit and none in the α -, γ -, or δ -subunits. Wild-type recovery was performed by suppressing at $\alpha 149GGGU$ with YFaFS_{ACCC}-W and adding either

Table 3. Wild-type recovery and UAA incorporation by frameshift suppression *in vivo*

mRNA	tRNA	EC ₅₀ (theoretical)*	n_H	n
$\alpha 149CGGG$	YFFS _{CCCG} -W [†]	56 \pm 2 (50) [‡]	1.8 \pm 0.07	8
$\alpha 149GGGU$	YFaFS _{ACCC} -W [†]	53 \pm 2 (50) [‡]	1.6 \pm 0.03	8
$\beta 9'$ GGGU	YFaFS _{ACCC} -Aba	16 \pm 0.9 (16) [‡]	1.3 \pm 0.08	7
$\delta 9'$ GGGU	YFaFS _{ACCC} -Nval	31 \pm 2 (36) [‡]	1.6 \pm 0.1	6
$\alpha 149CGGG$	YFFS _{CCCG} -WF1	190 \pm 3 (200) [§]	1.6 \pm 0.03	10

*EC₅₀ values from THG73-UAA incorporation by nonsense suppression.

[†]Rescue of wild-type recovery by frameshift suppression.

[‡]Ref. 14.

[§]Ref. 18.

wild-type or masked β mRNA to the injection mixture. Table 6, which is published as supporting information on the PNAS web site, shows the dramatic effect of masking one position on frameshift suppression. With a 1:1:1:1 ratio of α : β : γ : δ , the masked construct gives a 2.7-fold increase in I_{max} relative to wild type. As the amount of α -subunit (which contains the suppression site) is increased, the masking effect decreases to 1.5-fold and 1.2-fold with subunit ratios of 5:1:1:1 and 10:1:1:1, respectively. Calculations that assume two equally efficient quadruplet codons reproduce this trend (Table 6), suggesting that the $\alpha 149GGGU$ and the GGGU present in the β -subunit have similar suppression efficiencies.

Comparison of Frameshift and Nonsense Suppression Efficiencies. To compare frameshift and nonsense suppression, the $\alpha 149$ and $\beta 9'$ sites were studied in more detail. Suppression of $\alpha 149CGGG$ or GGGU with 10 ng of YFFS_{CCCG}-W or YFaFS_{ACCC}-W resulted in 38% and 48%, respectively, of the current from 10 ng of THG73-W suppression at $\alpha 149UAG$ (Table 4). Suppression of $\beta 9'$ UAG with 2 ng of THG73-L resulted in the largest I_{max} (Table 4). Suppression at $\beta 9'$ CGGG or GGGU with 2 ng of YFFS_{CCCG}-L or YFaFS_{ACCC}-L resulted in 14% and 36%, respectively, of the current from THG73-L (Table 4). We conclude that amber suppression is more efficient than frameshift suppression, in agreement with a trend previously seen in a eukaryotic cell-free translation system (16). In particular, the suppression efficiency observed here follows the order: THG73 > YFaFS_{ACCC} > YFFS_{CCCG}.

Interestingly, the yield of receptors from frameshift suppression at the $\beta 9'$ site was substantially improved by increasing the amount of tRNA injected. Suppression with 6 ng of YFFS_{CCCG}-L or

Table 4. Comparison of suppression efficiency, aminoacylation, and read-through *in vivo*

mRNA	tRNA	tRNA, ng	n	$I_{max} \pm SE^*$	% THG73 or % UAG
$\alpha 149UAG$	THG73-W	10	18	-4.8 \pm 2	100 [†]
$\alpha 149CGGG$	YFFS _{CCCG} -W	10	20	-1.8 \pm 0.3	38 [†]
$\alpha 149GGGU$	YFaFS _{ACCC} -W	10	13	-2.3 \pm 0.9	48 [†]
$\beta 9'$ UAG	THG73-L [‡]	2	15	-6.1 \pm 2	100 [†]
$\beta 9'$ CGGG	YFFS _{CCCG} -L [‡]	2	12	-0.84 \pm 0.2	14 [†]
$\beta 9'$ GGGU	YFaFS _{ACCC} -L [‡]	2	9	-2.2 \pm 0.5	36 [†]
$\beta 9'$ CGGG	YFFS _{CCCG} -L [‡]	6	13	-8.8 \pm 0.9	NA [†]
$\beta 9'$ GGGU	YFaFS _{ACCC} -L [‡]	6	13	-16 \pm 2	NA [†]
$\beta 9'$ UAG	THG73-dCA	2	13	-4.8 \pm 1	100 [†]
$\beta 9'$ CGGG	YFFS _{CCCG} -dCA	2	13	-0.42 \pm 0.8	8.8 [†]
$\beta 9'$ GGGU	YFaFS _{ACCC} -dCA	2	13	-0.092 \pm 0.02	1.9 [†]
$\beta 9'$ UAG	THG73-dCA	6	13	-8.2 \pm 1	100 [†]
$\beta 9'$ CGGG	YFFS _{CCCG} -dCA	6	12	-1.2 \pm 0.3	15 [†]
$\beta 9'$ GGGU	YFaFS _{ACCC} -dCA	6	11	-0.27 \pm 0.09	3.3 [†]
$\beta 9'$ UAG	—	—	13	-0.37 \pm 0.1	100 [§]
$\beta 9'$ CGGG	—	—	13	-0.085 \pm 0.03	23 [§]
$\beta 9'$ GGGU	—	—	13	-0.078 \pm 0.02	21 [§]

NA, not applicable; —, no tRNA.

*Average I_{max} (in μA) recorded at 1 mM ACh.

[†]THG73.

[‡]Currents in response to 10 μM and 1 mM ACh displayed a ratio of 0.1, as anticipated from the Hill equation fit for one wild-type receptor.

[§]UAG.

Table 5. Incorporation of two UAAs

α 149	tRNA	β or δ	tRNA	EC ₅₀ (theoretical)*	n _H	n
UAG	THG73-W	β 9'CGGG	YFFS _{CCCG} -Aba	14 ± 0.4 (16) [†]	1.7 ± 0.06	9
UAG	THG73-W	δ 9'GGGU	YFaFS _{ACCC} -Nval	41 ± 2 (36) [†]	1.9 ± 0.1	9
UAG	THG73-W	β 9'CGGG	YFFS _{CCCG} -L	50 ± 3 (50) [‡]	1.4 ± 0.08	20
UAG	THG73-WF1	β 9'CGGG	YFFS _{CCCG} -Aba	61 ± 3	1.5 ± 0.08	7
UAG	THG73-WF1	δ 9'GGGU	YFaFS _{ACCC} -Nval	180 ± 7	1.8 ± 0.1	6
UAG	THG73-WF1	β 9'CGGG	YFFS _{CCCG} -L	200 ± 7 (200) [‡]	1.3 ± 0.04	9

*EC₅₀ values from THG73-UAA incorporation by nonsense suppression.

[†]Ref. 14.

[‡]Ref. 18.

YFaFS_{ACCC}-L gave dramatic increases in I_{\max} , with a percentage change of 950% and 630%, respectively (Table 4). This large change in I_{\max} in response to a modest increase in tRNA concentration implicates a competition with endogenous triplet tRNA that responds nonlinearly to the amount of injected FS tRNA. A comparable increase in the amount of injected THG73-L led to complications due to reacylation of the tRNA by endogenous aaRSs (undesired) and incorporation of natural amino acids other than leucine, an issue that is addressed in detail in the following section and in *Discussion*.

Comparison of Aminoacylation of Suppressor tRNA and Read-Through of Suppression Sites. To evaluate aminoacylation *in vivo*, which is undesirable for any tRNA used to incorporate UAAs, the β 9' site was again studied, because most amino acids produce functional receptors when substituted at this position (14). In all experiments, tRNAs that had been ligated to dinucleotide deoxyCA (dCA) but did not contain an amino acid at the 3' end were injected to more closely mimic the biologically active, full-length tRNA. To maximize the potential for aminoacylation by endogenous aaRSs, 2-day incubations and relatively large mRNA injections (16.5 ng) were used. Surprisingly, THG73-dCA, which has been used extensively for UAA incorporation in *Xenopus* oocytes, showed significant aminoacylation *in vivo*, with an I_{\max} of -4.8 and -8.2 μ A for 2 and 6 ng of tRNA, respectively (Table 4). Note that under other conditions (less mRNA and shorter incubations), previous work has found no complications from aminoacylation using THG73-dCA in *Xenopus* oocytes (9, 14, 17). Still, the present results establish that THG73 is susceptible to aminoacylation by aaRSs, which is undesired. No aminoacylation was seen with 2 ng of THG73-L, suggesting that aminoacylation by endogenous aaRSs is more likely when nonaminoacylated THG73 is injected, as noted previously (17). Both FS tRNAs show much lower amounts of aminoacylation by aaRSs, as evidenced by the decrease in I_{\max} (Table 4). YFFS_{CCCG}-dCA shows only 8.8% and 15% of the I_{\max} of THG73-dCA at 2 and 6 ng, respectively. The most orthogonal suppressor is YFaFS_{ACCC}-dCA, with 1.9% and 3.3% of the I_{\max} of THG73-dCA at 2 and 6 ng, respectively. The orthogonality trend thus follows the order: YFaFS_{ACCC}-dCA > YFFS_{CCCG}-dCA > THG73-dCA. YFaFS_{ACCC} is the most orthogonal and efficient FS tRNA, and it therefore offers a viable replacement for THG73, especially when aminoacylation by aaRSs poses a problem *in vivo*.

Read-through at the β 9' site was also assessed by injection of mRNA only (Table 4). β 9'UAG showed the most read-through, presumably because there is only one in-frame stop codon before desired termination. β 9'CGGG and β 9'GGGU show 23% and 21% read-through, respectively, relative to the UAG stop codon. This finding is consistent with the idea that an endogenous triplet tRNA recognizing the first three bases of a quadruplet codon causes a -1 frameshift, which then presents multiple stop codons. Again, we designed this experiment to enhance signals from read-through by injecting large amounts of mRNA (50 ng). No current was detectable after injection of mRNA containing UAG,

CGGG, or GGGU at position α 149, confirming that this site is much less promiscuous than β 9'.

Incorporation of Two UAAs. To investigate the simultaneous incorporation of two UAAs, we again built on previous work that used THG73 to incorporate UAAs into the nAChR at positions α 149, β 9', and δ 9'. Importantly, EC₅₀ changes associated with mutations at these sites are independent of one another (18, 19), allowing one to qualitatively anticipate the consequences of multiple mutations. In particular, both β 9'Aba and δ 9'Nval produce predictable reductions in EC₅₀ that should be reproduced when combined with mutations at α 149 (14). That is, the previously noted 4-fold increase in EC₅₀ that is seen when the native tryptophan at α 149 is replaced by WF1 should persist when in combination with β 9'Aba or δ 9'Nval.

Successful incorporation of two UAAs to produce large ACh-induced currents could be seen when a 5-fold excess of mutant to wild-type mRNA was used. Suppression with α 149UAG/THG73-W and β 9'CGGG/YFFS_{CCCG}-L is a wild-type recovery experiment that gave the expected EC₅₀ for ACh of 50 μ M (Table 5). Maintaining β 9'CGGG/YFFS_{CCCG}-L but substituting α 149UAG/THG73-WF1 resulted in the anticipated 4-fold increase in EC₅₀ (Table 5) (18). For incorporation of two UAAs, α 149UAG/THG73-W or WF1 was combined with either β 9'CGGG/YFFS_{CCCG}-Aba or δ 9'GGGU/YFaFS_{ACCC}-Nval (Table 5 and Fig. 2; representative traces are shown in Fig. 4, which is published as supporting information on the PNAS web site). The α 149 WF1:W EC₅₀ ratios are 4.4 for the β 9' and δ 9' mutants. These experiments establish that frameshift suppression can be combined with nonsense suppression to incorporate two UAAs in a eukaryotic system.

Incorporation of Three UAAs. To demonstrate the incorporation of three UAAs, we combined the two-UAA incorporation experiments described above, taking advantage of the knowledge that




Fig. 2. Fits to the Hill equation (for incorporation of two UAAs). Data correspond to the rows in Table 5 as follows: row 1, open circles; row 2, open squares; row 4, filled circles; row 5, filled squares. Rows 3 and 6 are left out for clarity and have previously been reported (Table 3 and ref. 18).

Gene Construction and RNA Preparation. The α -, β -, γ -, and δ -subunits of nAChR were previously subcloned in the pAMV vector (25). All four in-frame CGGGs were mutated (shown in italics) to degenerate codons (*α 182CGC*, *β 234AGG*, *β 402AGG*, and *δ 1954AGG*), and one GGGT was mutated at the fourth position (*β 1AGC*); these constructs are termed “masked.” *α 149TAG*, CGGG, GGGT; *β 9'TAG*, CGGG, GGGT; and *δ 9'GGGT* mutations were placed on masked constructs by QuikChange site-directed mutagenesis (Stratagene). Mutations were verified by DNA sequencing (at the California Institute of Technology Sequencing/Structure Analysis Facility). Template DNA was linearized with NotI and mRNA prepared with the T7 mMessage mMachine kit. mRNA was purified by using the RNeasy Mini kit (Qiagen, Valencia, CA) and quantified by absorption at 260 nm.

THG73 and HSAS in pUC19 vector were previously made (10, 17). Genes for HSFSC_{CCCG}, THG73FS_{CCCG}, and YFFS_{CCCG} (sequence from ref. 6) with flanking EcoRI and BamHI overhangs were phosphorylated by using the Kinase Max kit, annealed, and ligated with T4 DNA ligase into EcoRI and BamHI linearized pUC19 vector as described in ref. 24. A73G; C2G, G3C, G4C; and C69G, C70G, G71C mutations (from ref. 7) were sequentially placed by QuikChange mutagenesis on the YFFS_{CCCG} construct to obtain YFaFS_{CCCG} (“a” refers to acceptor stem mutations). HSFSA_{CCCG} and YFaFSA_{CCCG} (sequence from ref. 7) were prepared by replacing the anticodon of HSFSC_{CCCG} and YFaFS_{CCCG} with ACCC by using QuikChange. All mutations were verified by DNA sequencing (at the California Institute of Technology Sequencing/Structure Analysis Facility). Template DNA for tRNA lacking the 3'CA was prepared by FokI digestion, and tRNA was transcribed by using the T7 MEGAshortscript kit. tRNA was desalted by using CHROMA SPIN-30 DEPC-H₂O columns (BD Biosciences), and concentration was determined by absorption at 260 nm.

dCA and dCA-UAA Ligation to Suppressor tRNA. dCA and 6-nitroveratryloxycarbonyl-protected dCA-UAA were coupled to suppressor tRNA by using T4 RNA ligase for 30 min as described in refs. 24 and 26, desalted by using CHROMA SPIN-30 DEPC-H₂O columns, and quantified by absorption at 260 nm. tRNA ligation efficiency was determined by MALDI mass spectrometry (26), and all tRNA dCA or dCA-UAA ligations were >75%.

In Vivo Suppression Experiments. Stage VI oocytes of *Xenopus laevis* were prepared as described in ref. 27. All tRNA were refolded at 65°C for 2 min, and tRNA-UAA were deprotected for 5 min by UV irradiation before injection (17). The injection volume for all experiments was 50 nl, and the incubation time was 44–52 h unless otherwise noted. Suppression of HSAS and HSFs (1.25 or 2.5 ng of tRNA) with 20 ng of mRNA in a subunit ratio of 2:5:1:1 for α : β 9'(UAG, CGGG, or GGGU): γ : δ was recorded after 1 or 2 days. Single UAA incorporation was performed by using 20–30 ng of

mRNA in a subunit ratio of 10:1:1:1 for α 149(CGGG or GGGU): β : γ : δ , 2:5:1:1 for α : β 9'GGGU: γ : δ , or 2:1:1:5 for α : β : γ : δ 9'GGGU with 4.8–16.5 ng of YFFS_{CCCG}/YFaFSA_{CCCG}-UAA. Comparison of β -masked and wild-type suppression contained 25 ng of total mRNA injected in the subunit ratio listed in Table 6 with 1:1 γ : δ and 10 ng of YFaFSA_{CCCG}-W. For comparison of suppression efficiency and aminoacylation of tRNA *in vivo*, all mRNA was normalized to the same concentration, and 16.5 ng of mRNA was injected in the subunit ratio 10:1:1:1 for α 149(UAG, CGGG, or GGGU): β : γ : δ or 2:5:1:1 for α : β 9'(UAG, CGGG, or GGGU): γ : δ with tRNA amounts listed in Table 4. For read-through experiments, 50 ng of mRNA in the ratio 2:5:1:1 for α : β 9'(UAG, CGGG, or GGGU): γ : δ was injected. Two UAA experiments were performed by injection of 20–30 ng of mRNA in a subunit ratio of 5:5:1:1 for α 149UAG: β 9'CGGG: γ : δ or 5:1:1:5 for α 149UAG: β : γ : δ 9'GGGU with 10–25 ng of each suppressor tRNA-UAA. For three UAAs, α 149W, β 9'Aba, and δ 9'Nval, 26 ng of mRNA in a ratio of 5:5:1:5 for α 149UAG: β 9'CGGG: γ : δ 9'GGGU was injected with 20 ng of each suppressor tRNA-UAA, and a second injection of 33 ng of each tRNA-UAA was done 24 h later. For α 149WF1, β 9'Aba, and δ 9'Nval, 8 ng of α 149UAG mRNA with 50 ng of THG73-WF1 was injected, and a second injection of 18 ng of mRNA with a subunit ratio of 5:1:5 for β 9'CGGG: γ : δ 9'GGGU with 25 ng of each YFFS_{CCCG}-Aba and YFaFSA_{CCCG}-Nval was performed 24 h later; oocytes were recorded 3 days after the first injection.

Electrophysiology. Recordings employed two-electrode voltage clamp on the OpusXpress 6000A (Axon Instruments, Union City, CA). ACh was stored at -20°C as a 1 M stock, diluted in Ca^{2+} -free ND96, and delivered to oocytes by computer-controlled perfusion system. For HSAS and HSFs experiments, the holding potential was -60 mV, and all UAA experiments were performed at either -60 or -80 mV. Dose–response data were obtained from at least nine ACh concentrations, and comparisons were tested at one drug concentration, except β 9'(UAG, CGGG, or GGGU) with tRNA-L, for which two concentrations, 10 μM and 1 mM, were used to check for aminoacylation (Table 4). Dose–response relations were fit to the Hill equation to determine EC_{50} and the Hill coefficient (n_{H}). All reported values are represented as a mean \pm SE of the tested oocytes [n is listed with each table].

Supporting Information. Masking experiments, representative traces for two UAA experiments (α 149W or WF1 and β 9'Aba), and oligonucleotides used in this study are detailed in Fig. 4 and Tables 6 and 7.

E.A.R. is a National Science Foundation Predoctoral Fellow. This work was supported by National Institutes of Health Grants NS-34407 and NS-11756.

- Dougherty, D. A. (2000) *Curr. Opin. Biotechnol.* **4**, 645–652.
- Beene, D. L., Dougherty, D. A., & Lester, H. A. (2003) *Curr. Opin. Neurobiol.* **13**, 264–270.
- Sakamoto, K., Hayashi, A., Sakamoto, A., Kiga, D., Nakayama, H., Soma, A., Kobayashi, T., Kitabatake, M., Takio, K., Saito, K., *et al.* (2002) *Nucleic Acids Res.* **30**, 4692–4699.
- Hino, N., Okazaki, Y., Kobayashi, T., Hayashi, A., Sakamoto, K., & Yokoyama, S. (2005) *Nat. Methods* **2**, 201–206.
- Zhang, Z., Alfonta, L., Tian, F., Bursulaya, B., Uryu, S., King, D. S., & Schultz, P. G. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 8882–8887.
- Hohsaka, T., Ashizuka, Y., Murakami, H., & Sisido, M. (1996) *J. Am. Chem. Soc.* **118**, 9778–9779.
- Hohsaka, T., Ashizuka, Y., Taira, H., Murakami, H., & Sisido, M. (2001) *Biochemistry* **40**, 11060–11064.
- Anderson, J. C., Wu, N., Santoro, S. W., Lakshman, V., King, D. S., & Schultz, P. G. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 7566–7571.
- Shafer, A. M., Kalai, T., Bin Liu, S. Q., Hideg, K., & Voss, J. C. (2004) *Biochemistry* **43**, 8470–8482.
- Monahan, S. L., Lester, H. A., & Dougherty, D. A. (2003) *Chem. Biol.* **10**, 573–580.
- Anderson, J. C., Magliery, T. J., & Schultz, P. G. (2002) *Chem. Biol.* **9**, 237–244.
- Saks, M. E., Sampson, J. R., & Abelson, J. N. (1994) *Science* **263**, 191–197.
- Murakami, H., Kourouklis, D., & Suga, H. (2003) *Chem. Biol.* **10**, 1077–1084.
- Kearney, P. C., Zhang, H. Y., Zhong, W., Dougherty, D. A., & Lester, H. A. (1996) *Neuron* **17**, 1221–1229.
- Grosjean, H., Droogmans, L., Giege, R., & Uhlenbeck, O. C. (1990) *Biochim. Biophys. Acta* **1050**, 267–273.
- Taira, H., Fukushima, M., Hohsaka, T., & Sisido, M. (2005) *J. Biosci. Bioeng.* **99**, 473–476.
- Saks, M. E., Sampson, J. R., Nowak, M. W., Kearney, P. C., Du, F. Y., Abelson, J. N., Lester, H. A., & Dougherty, D. A. (1996) *J. Biol. Chem.* **271**, 23169–23175.
- Zhong, W. G., Gallivan, J. P., Zhang, Y. O., Li, L. T., Lester, H. A., & Dougherty, D. A. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 12088–12093.
- Kearney, P. C., Nowak, M. W., Zhong, W., Silverman, S. K., Lester, H. A., & Dougherty, D. A. (1996) *Mol. Pharmacol.* **50**, 1401–1412.
- Labarca, C., Nowak, M. W., Zhang, H., Tang, L., Deshpande, P., & Lester, H. A. (1995) *Nature* **376**, 514–516.
- Nakamura, Y., Gojobori, T., & Ikemura, T. (2000) *Nucleic Acids Res.* **28**, 292.
- Magliery, T. J., Anderson, J. C., & Schultz, P. G. (2001) *J. Mol. Biol.* **307**, 755–769.
- England, P. M. (2004) *Biochemistry* **43**, 11623–11629.
- Nowak, M. W., Gallivan, J. P., Silverman, S. K., Labarca, C. G., Dougherty, D. A., & Lester, H. A. (1998) *Methods Enzymol.* **293**, 504–529.
- Nowak, M. W., Kearney, P. C., Sampson, J. R., Saks, M. E., Labarca, C. G., Silverman, S. K., Zhong, W., Thorson, J., Abelson, J. N., Davidson, N., *et al.* (1995) *Science* **268**, 439–442.
- Peterson, E. J., Shahgholi, M., Lester, H. A., & Dougherty, D. A. (2002) *RNA* **8**, 542–547.
- Quick, M. W., & Lester, H. A. (1994) in *Ion Channels of Excitable Cells*, ed. Narahashi, T. (Academic, San Diego), Vol. 19, pp. 261–279.